

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

UNCLASSIFIED

This Document
Reproduced From
Best Available Copy

DA 1179999

Armed Services Technical Information Agency

Reproduced by

DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address: Armed Services Technical Information Agency, Document Service Center, Knott Building, Dayton 2, Ohio.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.


UNCLASSIFIED

AD No 117999

437th FILE COPY

A SIMPLE PROOF AND SOME EXTENSIONS OF THE SAMPLING THEOREM

BY
EMANUEL PARZEN



TECHNICAL REPORT NO. 7

PREPARED UNDER CONTRACT Nonr-225 (21)
(NR-042-993)

FOR
OFFICE OF NAVAL RESEARCH

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

DECEMBER 22, 1956

A SIMPLE PROOF AND SOME EXTENSIONS
OF THE SAMPLING THEOREM

By
Emanuel Parzen

TECHNICAL REPORT NO. 7

PREPARED UNDER CONTRACT Nonr-225(21)
(NR-042-993)

FOR
OFFICE OF NAVAL RESEARCH

REPRODUCTION IN WHOLE OR IN PART IS PERMITTED FOR
ANY PURPOSE OF THE UNITED STATES GOVERNMENT

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

DECEMBER 22, 1956

A SIMPLE PROOF AND SOME EXTENSIONS
OF THE SAMPLING THEOREM

by
Emanuel Parzen

Abstract.

The sampling theorem states essentially that if the frequency spectrum, or Fourier transform, $g(\omega)$ of a time function $f(t)$ vanishes for ω outside some interval I , then $f(t)$ is completely determined by its values at certain discrete sampling points, whose density is proportional to the length of the interval I . This note gives a method of proof of the sampling theorem, both for the case where the interval I is centered at the origin and where it is not, which is somewhat simpler than the previously given proofs, and at the same time is more rigorous, and yields several useful generalizations to functions of several variables and random functions.

1. Introduction.

The sampling theorem, as first stated by Shannon [1], asserts essentially that if the frequency spectrum, or Fourier transform, $g(\omega)$ of a time function $f(t)$ vanishes for ω outside some interval I , then $f(t)$ is completely determined by its values at certain discrete sampling points, whose density is proportional to the length of the interval. The usual proof of the sampling theorem is based on expanding the Fourier transform $g(\omega)$ in a Fourier series. However, it appears that the result can be obtained more simply by expanding the kernel $e^{i\omega t}$ in a Fourier series, as a function of ω in I , which amounts to proving the sampling theorem for the functions of t given by $e^{i\omega t}$, with ω held fixed at some point in I . In this note, we use this

method to prove the sampling theorem. We also show how the method may be used to give simple proofs of various extensions of the sampling theorem, among them those due to Goldman [2], Kohlenberg [3], and Woodward and Davies [4]. The theorems we prove are the following.

Theorem I: Suppose that the function $f(t)$, defined for all real values of t , may be represented as a Fourier integral, for some positive constant W ,

$$(1.1) \quad f(t) = \int_{-2\pi W}^{2\pi W} e^{i\omega t} g(\omega) d\omega$$

where the function $g(\omega)$ may contain δ function terms, but not at $\omega = \pm 2\pi W$. Then, for every t ,

$$(1.2) \quad f(t) = \sum_{m=-\infty}^{\infty} f\left(\frac{m}{2W}\right) s\left(t - \frac{m}{2W}\right)$$

where

$$(1.3) \quad s(t) = \frac{1}{4\pi W} \int_{-2\pi W}^{2\pi W} e^{i\omega t} d\omega$$

$$(1.4) \quad = \frac{\sin 2\pi W t}{2\pi W t}$$

Theorem II: Suppose that $f(t_1, \dots, t_K)$, defined as a function of K variables, may be represented as a Fourier integral, for positive W_1, \dots, W_K ,

$$(1.5) \quad f(t_1, \dots, t_K) = \int_{-2\pi W_1}^{2\pi W_1} \dots \int_{-2\pi W_K}^{2\pi W_K} d\omega_1 \dots d\omega_K e^{i(\omega_1 t_1 + \dots + \omega_K t_K)} g(\omega_1, \dots, \omega_K)$$

where $g(\omega_1, \dots, \omega_K)$ may contain δ function terms, but not at the 2^K points of the form $(\pm 2\pi W_1, \dots, \pm 2\pi W_K)$. Then

$$(1.6) \quad f(t_1, \dots, t_K) = \sum_{m_1, \dots, m_K = -\infty}^{\infty} f\left(\frac{m_1}{2W_1}, \dots, \frac{m_K}{2W_K}\right) s\left(t_1 - \frac{m_K}{2W_K}\right) \dots s\left(t_K - \frac{m_K}{2W_K}\right)$$

Theorem III: Let $f(t)$ be a stationary random function (as defined by Doob [5]) whose covariance (or autocorrelation) function $R(\tau) = E f(t) f(t+\tau)$ may be written,

$$(1.7) \quad R(\tau) = \int_{-2\pi W}^{2\pi W} e^{i\omega\tau} G(\omega) d\omega$$

where $G(\omega)$ is the power spectrum of the function, which may contain δ function terms, but not at $\omega = \pm 2\pi W$. Then

$$(1.8) \quad f(t) = \sum_{m=-\infty}^{\infty} f\left(\frac{m}{2W}\right) s\left(t - \frac{m}{2W}\right)$$

where the infinite series in (18) converges stochastically (in the sense of convergence in mean square).

Theorem IV: Suppose that $f(t)$ may be represented as a Fourier integral, for positive W_0 and W ,

$$(1.9) \quad f(t) = \int_{2\pi W_0}^{2\pi(W_0 + W)} e^{i\omega t} g(\omega) d\omega + \int_{-2\pi(W_0 + W)}^{-2\pi W_0} e^{i\omega t} g(\omega) d\omega$$

where the function $g(\omega)$ may possess δ function terms, but not at $\omega = \pm 2\pi W_0$ or $\omega = \pm 2\pi(W_0 + W)$ or $\omega = \pm 2\pi(rW - W_0)$ where r is the integer such that

$$(1.10) \quad 2 \frac{W_0}{W} \leq r < 2 \frac{W_0}{W} + 1$$

Let

$$(1.11) \quad \begin{aligned} s(t) = & \frac{1}{2\pi W} \int_{2\pi W_0}^{2\pi(rW + W_0)} d\omega [\cos \omega t + \sin \omega t \cot \pi r W k] \\ & + \frac{1}{2\pi W} \int_{2\pi(rW - W_0)}^{2\pi(W_0 + W)} d\omega [\cos \omega t + \sin \omega t \cot \pi(r + 1) W k] \end{aligned}$$

where k is a constant satisfying the condition that rWk and $(r + 1)Wk$ are not equal to any of the integers $0, \pm 1, \dots$. Then, for every t ,

$$(1.12) \quad f(t) = \sum_{m=-\infty}^{\infty} f\left(\frac{m}{W}\right) s\left(t - \frac{m}{W}\right) + f\left(\frac{m}{W} + k\right) s\left(\frac{m}{W} + k - t\right)$$

If W_0/W is an integer, then with $k = 1/2W$, (1.12) reduces to (1.4).

It will be clear from what follows that Theorems II and III can be extended in the same way that Theorem I is extended by Theorem IV.

2. Proofs of Theorems I-III.

We start with the following basic fact from the theory of Fourier series. For any real number t , and ω such that $|\omega| < 2\pi W$,

$$(2.1) \quad e^{i\omega t} = \sum_{m=-\infty}^{\infty} e^{i\omega \frac{m}{2W}} s\left(t - \frac{m}{2W}\right) .$$

The infinite series in (2.1) converges in general for $|\omega| < 2\pi W$, but not for $\omega = \pm 2\pi W$. However, the consecutive sums

$$S_M(\omega, t) = \sum_{|m| \leq M} e^{i\omega \frac{m}{2W}} s\left(t - \frac{m}{2W}\right)$$

are, for fixed t , uniformly bounded for $|\omega| \leq 2\pi W$; that is, there is a constant K such that $|S_M(\omega, t)| \leq K$ for $M = 0, 1, 2, \dots$ and $|\omega| \leq 2\pi W$.

A not quite rigorous way of verifying (2.1) is as follows. The right hand side of (2.1) may be written

$$(2.2) \quad \frac{1}{4\pi W} \int_{-2\pi W}^{2\pi W} d\lambda e^{i\lambda t} \sum_{m=-\infty}^{\infty} e^{i(\omega-\lambda)\frac{m}{2W}} .$$

It may be verified that

$$(2.3) \quad \sum_{m=-\infty}^{\infty} e^{im\alpha} = 2\pi \sum_{n=-\infty}^{\infty} \delta(\alpha + 2\pi n)$$

and, for any function $f(x)$,

$$(2.4) \quad \int_I f(x) \delta\left(\frac{x-a}{c}\right) dx = c f(a)$$

if the point a lies in the interval of integration I ; otherwise the integral in (2.4) is zero. Now, for $|\omega| < 2\pi W$, the only value of n such that $|\omega + 4\pi n W| \leq 2\pi W$ is $n = 0$. Consequently, in view of (2.3) and (2.4),

it is seen that the value of (2.2) is $e^{i\omega t}$, for $|\omega| < 2\pi W$, which verifies (2.1).

Next, to prove Theorems I and II, merely replace $e^{i\omega t}$ in (1.1) and (1.5) by the infinite series of (2.1); that the order of integration and summation may be interchanged follows rigorously by the theory of Lebesgue integration.

Next, to prove Theorem III, we first note the fact that, from (1.7) it follows that $f(t)$ may be represented as a stochastic integral as follows:

$$(2.6) \quad f(t) = \int_{-2\pi W}^{2\pi W} e^{i\omega t} dZ(\omega) \quad .$$

By the theory of stochastic integrals, (1.8) follows from (2.6) if one shows that

$$\lim_{M \rightarrow \infty} \int_{-2\pi W}^{2\pi W} \left| e^{i\omega t} - S_M(\omega, t) \right|^2 G(\omega) d\omega = 0$$

which follows from the facts stated at the beginning of this section.

3. Proof of Theorem IV.

To prove (1.12) we begin by conjecturing that for ω such that $2\pi W_0 < |\omega| < 2\pi(W_0 + W)$, $e^{i\omega t}$ may be written in the form

$$(3.1) \quad e^{i\omega t} = \sum_{m=-\infty}^{\infty} e^{i\omega \frac{m}{W}} s_1\left(t - \frac{m}{W}\right) + \sum_{m=-\infty}^{\infty} e^{i\omega\left(\frac{m}{W} + k\right)} s_2\left(t - \frac{m}{W} - k\right)$$

for some constant k , and functions $s_1(t)$ and $s_2(t)$ which are of the

form, for $i = 1, 2$,

$$(3.2) \quad s_i(t) = \int_I e^{it\omega} S_1(\omega) d\omega$$

where the region of integration I consists of the intervals $(2\pi W_0, 2\pi(W_0 + W))$ and $(-2\pi(W_0 + W), -2\pi W_0)$. In view of (3.2), the right hand side of (3.1) may be written

$$(3.3) \quad \int_I d\lambda S_1(\lambda) e^{i\lambda t} \sum_m e^{i(\omega-\lambda)\frac{m}{W}} \\ + \int_I d\lambda S_2(\lambda) e^{i\lambda t} e^{ik(\omega-\lambda)} \sum_m e^{i(\omega-\lambda)\frac{m}{W}}.$$

We next evaluate (3.3) by means of (2.3) and (2.4).

It is readily verified that, for ω in I , $\omega + 2\pi nW$ lies in I only for $n = 0$, and for one other value of n , which we denote by $N(\omega)$, whose values are given in Table A in terms of the integer r given by (1.10).

Table A

| For ω in the interval | $N(\omega) =$ | $\omega + 2\pi N(\omega)W$ lies in the interval |
|--|---------------|---|
| $J_1 : 2\pi W_0 < \omega < 2\pi(rW - W_0)$ | $-r$ | $-J_1$ |
| $-J_1 : -2\pi(rW - W_0) < \omega < -2\pi W_0$ | r | J_1 |
| $J_2 : 2\pi(rW - W_0) < \omega < 2\pi(W_0 + W)$ | $-(r+1)$ | $-J_2$ |
| $-J_2 : -2\pi(W_0 + W) < \omega < -2\pi(rW - W_0)$ | $r+1$ | J_2 |

Then, for ω in I , the value of (3.3) is given by

$$(3.4) \quad 2\pi W e^{i\omega t} [S_1(\omega) + S_2(\omega)] + 2\pi W e^{it(\omega + 2\pi N(\omega)W)}$$

$$[S_1(\omega + 2\pi N(\omega)W) + e^{-i2\pi N(\omega)Wk} S_2(\omega + 2\pi N(\omega)W)]$$

Now, let $S_1(\omega)$ and $S_2(\omega)$ be defined so that they vanish for ω outside I , and for $\omega = \pm 2\pi(rW - W_0)$, and for other values of ω in I satisfy the relations

$$(3.5) \quad 2\pi W S_1(\omega) = 1 - 2\pi W S_2(\omega), \quad 2\pi W S_2(\omega) = [1 - e^{i2\pi N(\omega)Wk}]^{-1}$$

where we now choose k so that rWk and $(r+1)Wk$ are not equal to any of the integers $0, \pm 1, \pm 2, \dots$. In view of Table A it may be verified that the following relations hold for ω in I , where an asterisk denotes a complex conjugate:

$$N(\omega) = -N(-\omega), \quad S_1(\omega) = S_2(-\omega)$$

$$S_2(\omega + 2\pi N(\omega)W) = S_2^*(\omega) = S_2(-\omega),$$

$$S_1(\omega + 2\pi N(\omega)W) = S_1^*(\omega) = S_1(-\omega),$$

$$S_1(\omega + 2\pi N(\omega)W) + e^{-i2\pi N(\omega)Wk} S_2(\omega + 2\pi N(\omega)W)$$

$$= [S_1(\omega) + e^{i2\pi N(\omega)Wk} S_2(\omega)]^* = 0,$$

$$2\pi W(S_1(\omega) + S_2(\omega)) = 1$$

In view of these relations, the value of (3.4) is $e^{i\omega t}$ for ω in I , except perhaps for $\omega = \pm 2\pi(rW - W_0)$, and (3.1) holds with $s_1(t)$ and $s_2(t)$ given by (3.2) and $S_1(\omega)$ and $S_2(\omega)$ given by (3.5).

We now write (3.1) in a more convenient form. Since $S_1(\omega) = S_2(-\omega)$, it follows that $s_1(t) = s_2(-t)$. Define $s(t) = s_1(t)$. Then, since $S_1^*(\omega) = S_1(-\omega)$,

$$(3.6) \quad s(t) = \int_I e^{i\omega t} S_1(\omega) d\omega = 2 \int_{+2\pi W_0}^{2\pi(W_0 + W)} \text{Real} [e^{i\omega t} S_1(\omega)] d\omega .$$

Now

$$(3.7) \quad 2 \text{Real} [e^{i\omega t} S_1(\omega)] = \frac{1}{2\pi W} \left[\frac{\cos \omega t - \cos (\omega t + 2\pi N(\omega) W k)}{1 - \cos 2\pi N(\omega) W k} \right] \\ = \frac{1}{2\pi W} \left[\cos \omega t + \sin \omega t \cot \pi N(\omega) W k \right] .$$

Thus (1.12) has been established for the complex exponentials $e^{i\omega t}$ for $2\pi W_0 < |\omega| < 2\pi(W_0 + W)$, with the possible exception of $\omega = \pm 2\pi(rW - W_0)$. To establish (1.12) for any function $f(t)$ satisfying (1.9), merely replace $e^{i\omega t}$ in (1.9) by the infinite series representing it.

REFERENCES

- [1] C. E. Shannon, Proc. Inst. Radio Engrs., 37 (1949), p. 10.
- [2] S. Goldman, Information Theory, New York, 1953.
- [3] A. Kohlenberg, J. Appl. Physics, 24 (1953), p. 1432.
- [4] P. M. Woodward, Probability and Information Theory, London, 1953.
- [5] J. L. Doob, Stochastic Processes, New York, 1953, p. 95 and p. 473.

Acknowledgement. It is a pleasure to thank Professor L. M. Zadeh for several helpful conversations.

STANFORD UNIVERSITY

Technical Reports Distribution List Contract Nonr-225(21) (NR-042-993)

| | | | |
|--|---|--|---|
| Chief, Bureau of Ships Dept. of the Navy Washington 25, D. C. Attn: H. Weingarten Code 223 | 1 | Scientific Section Office of Naval Research Dept. of the Navy 1000 Geary St. San Francisco 9, Calif. | 2 |
| Commanding Officer U. S. Naval Powder Factory Indianhead, Md. Attn: F. Freshman, R. and D. | 1 | Statistical Engineering Lab. National Bureau of Standards Washington 25, D. C. | 1 |
| Director, Naval Research Lab. Washington 25, D. C. Attn: Technical Information Officer | 6 | Dept. of Statistics University of California Berkeley 4, Calif. | 1 |
| Document Service Center A.S.T.I.A. Knott Building Dayton 2, Ohio Attn: DSC-S | 5 | Statistical Laboratory University of Washington Seattle 5, Wash. | 1 |
| Dept. of Mathematical Statistics University of North Carolina Chapel Hill, N. C. | 1 | Professor O. P. Aggarwal Statistical Lab. Purdue University Lafayette, Indiana | 1 |
| Office of Naval Research Dept. of the Navy 17th and Constitution Aves. Washington 25, D. C. Attn: Code 433 | 1 | Dr. Stephen G. Allen 244 Laurel St. San Francisco, Calif. | 1 |
| Office of the Asst. Naval Attache for Research Naval Attache American Embassy Navy No. 100 Fleet Post Office New York, N. Y. | 2 | Professor Fred C. Andrews Mathematics Dept. University of Nebraska Lincoln 8, Nebr. | 1 |
| Office of Technical Services Dept. of Commerce Washington 25, D. C. | 1 | Professor T. W. Anderson Dept. of Mathematical Statistics Columbia University New York 27, N. Y. | 1 |
| | | Professor David Blackwell Dept. of Statistics University of California Berkeley 4, Calif. | 1 |
| | | Dr. Julius R. Blum Dept. of Mathematics Indiana University Bloomington, Indiana | 1 |

| | | | |
|--|---|---|---|
| Dr. Paul Blunk Box 532 Fair Oaks, Calif. | 1 | Professor E. J. Gumbel Industrial Engr. Dept. 409 Engineering Bldg. Columbia University New York 27, N. Y. | 1 |
| Dr. Charles Boll Hughes Aircraft Co. Bldg. 12, Room 2343 Culver City, Calif. | 1 | Professor Leonid Hurwicz School of Business Administration University of Minnesota Minneapolis 14, Minn. | 1 |
| Professor W. G. Cochran Biostatistics Dept. The Johns Hopkins University Baltimore 5, Md. | 1 | Professor Stanley Isaacson 716 Eastmont Dr., Apt. 2C Greensburg, Pa. | 1 |
| Professor Lee Cronbach Bureau of Education Research 1007 S. Wright Champaign, Ill. | 1 | Professor Leo Katz Mathematics Dept. Michigan State University East Lansing, Mich. | 1 |
| Dr. Joseph Daly Bureau of the Census Washington 25, D. C. | 1 | Professor Tosio Kitagawa Mathematical Institute Faculty of Science Kyusyu University Fukuoka, Japan | 1 |
| Dr. Francis Dresch Stanford Research Institute 1915 University Ave. Palo Alto, Calif. | 1 | Dr. Dennis V. Lindley Statistical Lab. University of Cambridge Cambridge, England c/o Office of Naval Research London Branch Navy No. 100, Fleet Post Office New York, N. Y. | 1 |
| Professor Meyer Dwass Dept. of Mathematics Northwestern University Evanston, Ill. | 1 | Mr. Monroe Norden Research Division Engineering Statistics Group College of Engineering New York University 401 W. 205 St. New York 54, N. Y. | 1 |
| Mr. Murray A. Geisler Logistics Section The RAND Corporation 1700 Main St. Santa Monica, Calif. | 1 | Professor Stanley Reiter Dept. of Economics Furdue University Lafayette, Indiana | 1 |
| Mr. Geoffrey Gregory 4 Osborne Grove Gatley, Cheadle Cheshire, England c/o Office of Naval Research London Branch Navy No. 100, Fleet Post Office New York, N. Y. | 1 | | |

Dr. A. R. Roy
Statistical Wing
Indian Council of Agricultural
Research
Linlithgow Ave.
New Delhi, India

1

Professor Herman Rubin
Dept. of Mathematics
University of Oregon
Eugene, Oregon

1

Dr. Jagdish Rustagi
Dept. of Mathematics
College of Engineering and Science
Carnegie Inst. of Tech.
Pittsburgh 13, Pa.

1

Professor Seymour Sherman
Moore Sch. of Electrical Engineering
University of Pennsylvania
Philadelphia 4, Pa.

1

Dr. Milton Sobel
Bell Telephone Labs.
555 Union Blvd.
Allentown, Pa.

1

Professor Herbert Solomon
Teachers' College
Columbia University
New York 27, N. Y.

1

Dr. Dan Teichroew
Numerical Analysis Research
405 Hilgard Ave.
Los Angeles 24, Calif.

1

Professor Donald Truax
Mathematics Dept.
University of Kansas
Lawrence, Kansas

1

Mr. Cesareo Villegas
Institute de Matematica y
Estadistica
Av. J. Herrera y Reissig 565
Montevideo, Uruguay

1

Professor W. Allen Wallis
Committee on Statistics
University of Chicago
Chicago 37, Ill.

1

Professor Irving Weiss
Bell Telephone Labs.
1600 Osgood St.
North Andover, Mass.

1

Dr. Oscar Wesler
Dept. of Mathematics
University of Michigan
Ann Arbor, Mich.

1

Dr. John D. Wilkes
Office of Naval Research
Code 200
Washington 25, D. C.

1

Professor S. S. Wilks
Room 110, Fine Hall
Box 708
Princeton, N. J.

1

Professor J. Wolfowitz
Mathematics Dept.
Cornell University
Ithaca, N. Y.

1

Professor M. A. Woodbury
Dept. of Mathematics
New York University
New York 53, N. Y.

1

Dr. Eugene Lukacs
Dept. of Mathematics
Catholic University
Washington 17, D. C.

1

Additional copies for project
leader and assistants and
reserve for future requirements 50

AD
1179999

A Mathematical Restatement of Hypotheses in
 "A SIMPLE PROOF AND SOME EXTENSIONS
 OF THE SAMPLING THEOREM"^{1/}

by

Emanuel Parzen

This report was written in a terminology which would presumably be most readable to communication engineers. For the reader interested in mathematical precision, the following restatements of hypotheses should be noted.

In Theorem I, for (1.1) read

$$(1.1) \quad f(t) = \int_{-2\pi W}^{2\pi W} e^{i\omega t} dV(\omega)$$

where $V(\omega)$ is a function of bounded variation continuous at $\omega = \pm 2\pi W$.

In Theorem II, for (1.5) read

$$(1.5) \quad f(t_1, \dots, t_K) = \int_{-2\pi W_1}^{2\pi W_1} \dots \int_{-2\pi W_K}^{2\pi W_K} e^{i(\omega_1 t_1 + \dots + \omega_K t_K)} dV(\omega_1, \dots, \omega_K)$$

where $V(\omega_1, \dots, \omega_K)$ is a function of bounded variation which assigns measure zero to the set $\{(\omega_1, \dots, \omega_K) : \omega_i = \pm 2\pi W_i \text{ for some } i\}$.

In Theorem III, for (1.7) read

$$(1.7) \quad R(\tau) = \int_{-2\pi W}^{2\pi W} e^{i\omega \tau} dF(\omega)$$

^{1/} Technical Report No. 7, Department of Statistics, Stanford University, Stanford, California. Prepared under Office of Naval Research Contract Nonr-225(21).

where the spectral distribution function $F(\omega)$ is monotone non-decreasing and continuous at $\omega = \pm 2\pi W$.

In Theorem IV, for (1.9) read

$$(1.9) \quad f(t) = \int_{2\pi W_0}^{2\pi(W_0 + W)} e^{i\omega t} dV(\omega) + \int_{-2\pi(W_0 + W)}^{-2\pi W_0} e^{i\omega t} dV(\omega)$$

where $V(\omega)$ is a function of bounded variation continuous at

$\omega = \pm 2\pi W_0$, $\omega = \pm 2\pi(W_0 + W)$, and $\omega = \pm 2\pi(rW - W_0)$, where r is defined by (1.10).